

# BIOCHEMICAL ANALYSIS OF FLOWERS OF *Vinca major*, A MEDICINAL WEED PLANT OF HILLY AREAS OF PAKISTAN

\*Arshad Javaid<sup>1</sup>, Malik F. H. Ferdosi<sup>2</sup>, Iqra Haider Khan<sup>1</sup>, Amna Shoaib<sup>1</sup>, Hafiz M. Saeed<sup>2</sup>, Muhammad Abrar Ul Hassan<sup>1</sup>

<sup>1</sup>Department of Plant Pathology, Faculty of Agricultural Sciences, University of the Punjab, Lahore, Pakistan

<sup>2</sup>Department of Horticulture, Faculty of Agricultural Sciences, University of the Punjab, Lahore, Pakistan

\*Corresponding author's email: [arshad.iags@pu.edu.pk](mailto:arshad.iags@pu.edu.pk), [arshadjpk@yahoo.com](mailto:arshadjpk@yahoo.com)

## ABSTRACT

*Vinca major* L. is an evergreen perennial weed of family Apocynaceae, growing naturally in hilly areas of Hazara region of Khyber Pakhtunkhwa, Pakistan. Flowers of this weed were collected from Khanspur, Ayubia during June 2021, shade dried and extracted in pure methanol for two weeks. After filtration, the extract was analyzed by GC-MS for the identification of possible bioactive compounds.  $\alpha$ -Amyrin was the major compound (32.49%) followed by lup-20(29)-en-3-ol, acetate, (3 $\beta$ )- (25.72%). Moderately abundant compounds included  $\gamma$ -sitosterol (8.78%),  $\beta$ -amyron (7.25%), cyclohexane, 1,3,5-triphenyl- (7.01%), olean-12-en-3-ol, acetate, (3 $\beta$ )- (5.68%), and *n*-hexadecanoic acid (3.18%). Some of the identified compounds from *V. major* have various important biological properties including anti-inflammatory, antimicrobial, anticancer, antimalarial, antioxidant, antidiabetic, antitumor, etc.

**Keywords:** Bioactive compounds, Flower extract, GC-MS analysis, Khyber Pakhtunkhwa, *Vinca major*.

## INTRODUCTION

Indigenous medicinal plants represent a reservoir of natural products that have been used by human beings since ancient times. This has resulted in the use of large number of plants with curative properties against several diseases in many parts of the world (Dar *et al.*, 2017; Jamshidi-Kia *et al.*, 2018; Khan and Javaid, 2020, 2021). Cost and side effects of conventional medications are the major factors responsible for revival of herbal medicines (Uritu *et al.*, 2018). Recent literature also shows the prevalent use of herbs as medicines across the globe (Saravanan *et al.*, 2018; Khan and Javaid, 2019). So far, a lot of important drugs are derived from traditional medicinal herbs (Manandhar *et al.*, 2019; Salmerón-Manzano *et al.*, 2020). A variety of compounds comprised of alkaloids, tannins, steroids, saponins, phenolic acids, terpenoids and quinones have been identified, which possess cardioprotective, anticancer, antioxidant, antimicrobial, anti-insect and a variety of other properties (Tungmunnithum *et al.*, 2018).

Comment [h1]: Write TITLE in upper case.

Comment [h2]: Write scientific names of organisms in lower case/sentence format.

Comment [h3]: Mention corresponding author with a \*. Only one author should be a correspondent.

Comment [h4]: 1-Abstract up to 250 words, in a single paragraph.  
2- All abbreviations should be explained at their citations. Common abbreviations such as GC-MS, ANOVA, LSD etc do not need explanation.  
3-Avoid too many abbreviations for smooth reading of the manuscript.  
4-Only use an abbreviation if it is to be used at certain positions in the subsequent text.

Comment [h5]: Full scientific names of the organisms used in the study should be followed by authority at their first citation.

Comment [h6]: No space between a digit and %.

Comment [h7]: Once writing the full scientific name at their first citation in the text, genus name should be abbreviated at all subsequent positions in the text.

Comment [h8]: Last sentence of the Abstract should be a conclusion of the study.

Comment [h9]: Up to 6 keywords in alphabetical order.

Comment [h10]: Introduction should be brief, composed of 2 to 3 paragraphs. Latest references up to the current year should be cited.

Comment [h11]: Follow this format for citing references in the text. Arrange references in the text in chronological order.

Pakistan is enriched in medicinal herbs because of varied climatic conditions (Ullah, 2017; Javed *et al.*, 2021). Many of these medicinal plants have been explored for their antifungal, antibacterial, herbicidal and other bioactive properties (Javaid *et al.*, 2020, 2021a; Banaras *et al.*, 2021; Ferdosi *et al.*, 2021a). These are scattered over a large area and about 600 species known well in Pakistan for their medicinal values (Shinwari, 2010). *Vinca major* of family Apocynaceae, is an evergreen perennial weed plant that has medicinal and ornamental values (Arora *et al.*, 2010). It grows well in full sun as well as in shade with a height of up to 25 cm while spreading indefinitely. Phytochemical screening of *V. major* revealed the presence of saponins, alkaloids, phenols, organic acids and sterols exhibiting antioxidant, antimicrobial, antidiabetic and hypotensive properties (Singh *et al.*, 2014; Wei and Liu, 2021). *Vinca* extracts have diverse compounds that act as antineoplastic agents and are also used to treat choriocarcinoma, lymphomas and hodgkin's diseases (Vishwakarma and Prajapati, 2019; González-Burgos and Gómez-Serranillos, 2021). Moreover, aerial plant parts are used traditionally to treat diarrhea, piles, leukemia, malaria, diabetes, sore throat, cough and diabetes (Rajput *et al.*, 2011; Ajaib *et al.*, 2014; Comfort *et al.*, 2019). However, studies regarding phytochemical analysis of flowers of this plant from Pakistan are scarce. Thus, the present study was conducted to identify phytoconstituents of *V. major* flowers through GC-MS analysis and to document its medicinal importance.

**Comment [h12]:** Differentiate with a, b, c if an author has more than one papers in the same year.

**Comment [h13]:** Give a hypothesis of mention what was missing that you have done in the present study.


**Comment [h14]:** Last sentence of introduction should be the objectives of your study.

## MATERIALS AND METHODS

Disease free and mature flowers of *V. major* were collected from Murree, Pakistan. Plucked flowers were packed in plastic bags and shifted to laboratory for further analysis. Flowers were washed in water and completely dried at 35 °C in a hot air oven. After evaporating moisture from flower, the dried flowers were then finely crushed into a powdered form with pestle and mortar. Ten grams of this material were soaked in 50 mL of analytical graded methanol and kept for two weeks so that maximum bioactive compounds can be extracted. Thereafter, the extract was filtered through a filter paper. Following filtration, 2 mL extract was collected in a 5 mL vial for GC-MS analysis.

**Comment [h15]:** 1-Methodology can be divided in to headings as per requirement of the paper.  
2- Write Methodology in such a complete way that another person can repeat it.  
3-Use SI units/MKS system.  
4-Write mg L<sup>-1</sup> instead of mg/L.  
5-Write mL and µL instead of ml and µl, respectively.

**Comment [h16]:** Put a single space between a digit and its unit except %.



**Fig. 1:** *Vinca major* growing in Murree, Pakistan.

**Comment [h17]:** Place all Figures and Tables at their appropriate positions in the text.

Various biologically active compounds were identified from methanolic flower extract using GC-MS analysis as described by Ferdosi *et al.* (2020). Gas chromatography (GC) was performed on 7890B Model Machine, Agilent Technologies (USA) with Column DB-5ms having dimensions of 30 m × 0.25 μm × 0.25 μm. Helium, an inert gas (having 4 amu atomic mass) was used as a carrier gas. Injection volume was 1 μL, while oven initial temperature was 80 °C that was raised to 10 °C min<sup>-1</sup> up to 300 °C. MS analysis was performed on 5977A Model Machine, Agilent Technologies (USA) with scan range of 50–500 m/z; solvent delay time was 5 min. Source temperature was 230 °C with 50 min run time. The resulted spectrum was then analyzed with NIST library of 2017 version for the phytochemical characterization and the compounds were arranged in the ascending order of their retention times. Relative abundance of the compounds was analyzed using of peaks heights in the chromatogram. Chemical structures of various compounds in the extract were drawn using ChemDraw software.

In order to collect information on biological activities of the identified compounds, different databases including Science Direct, PubMed, SciELO, Google Scholar, Directory of Open Access Journals and Crossref were surveyed.

## RESULTS AND DISCUSSION


Fourteen compounds were detected in GC-MS analysis of methanolic flower extract of *V. major* (Fig. 2). Details of the identified compounds are presented in Table 1. The most abundant compound was α-amyrin (32.49%) followed by lup-20(29)-en-3-ol, acetate, (3β)- (25.72%). Four compounds namely γ-sitosterol (8.78%), β-amyrone (7.25%), cyclohexane, 1,3,5-triphenyl- (7.01%), olean-12-en-3-ol, acetate, (3β)- (5.68%) and *n*-hexadecanoic acid (3.18%) were ranked as moderately abundant compounds. Compounds such as campesterol (2.11%), cyclododecane (1.17%), pentadecanoic acid, 14-methyl-, methyl ester (1.32%), heneicosane (1.31%), 2-ethylacridine (1.23%), 1,2-bis(trimethylsilyl)benzene (1.42%), were categorized as less abundant ones. Structures of major compounds are shown in Fig. 3.

The principal compound in the flower extract was α-amyrin. Earlier, this compound has been reported in *Cirsium arvense* (Ferdosi *et al.*, 2021b), *Myrcianthes pungens* (Cardoso *et al.*, 2020) and *Strobilanthes callosus* (Singh *et al.*, 2002). It is known for its antimicrobial, antioxidant and anti-inflammatory properties (Singh *et al.*, 2002). In addition, it also showed inhibitory effects against human oxidosqualene cyclase (Chen *et al.*, 2017). Its pharmacological activity is also known in gastrointestinal tract and immunological system (Nogueira *et al.*, 2019), and in the treatment of gingivitis and periodontitis (Pinto *et al.*, 2008). Likewise, β-amyrone isolated from oil-resins of *Protium paniculatum* also showed anti-inflammatory activity (de Almeida *et al.*, 2015).

The second most abundant compound lup-20(29)-en-3-ol, acetate, (3β)-, also known as lupeyl acetate and lupeol acetate, belongs to triterpenoids group. This compound was isolated from bark of *Artocarpus integra* and showed anticancer activity against breast cancer cells MCF-7 (Suwito *et al.*, 2016). It also possesses numerous other biological activities including anti-inflammatory, antituberculosis, antimalarial, antimicrobial and antinociceptive (Prachayasittikul *et al.*, 2010; Chen *et al.*, 2012). An isomer of this compound, olean-12-en-3-ol, acetate, (3β)- also known as β-amyrin 3- acetate, was found as a moderately occurring compound in this study. It

**Comment [h18]:** 1-Results and Discussion can be write separately or combined.  
2-Discussion should not be like that of Introduction.  
3-Instead of just citing previous literature, first give a brief account of a salient finding and then discuss it in the light of available literature especially the recent one.

possesses anti-inflammatory potential and also inhibits growth of *Staphylococcus aureus* (Hichri *et al.*, 2003; Akihisa *et al.*, 2010). It has been found as a major constituent in flowers of *C. arvense* and (Fernandes *et al.*, 2013; Ferdosi *et al.*, 2021b).




**Fig. 1:** GC-MS chromatogram of methanolic flower extract of *Vinca major*.

**Table 1:** Compounds identified in methanolic flower extract of *Vinca major* through GC-MS analysis.

Sr. No.	Names of compounds	Molecular formula	Molecular weight	Retention time (min)	Peak area (%)
1	5-Dodecene, (E)-	C <sub>12</sub> H <sub>24</sub>	168.31	9.366	1.17
2	Pentadecanoic acid, 14-methyl-, methyl ester	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	270.45	14.367	1.32
3	<i>n</i> -Hexadecanoic acid	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	256.42	14.768	3.18
4	9,12,15-octadecatrienoic acid, methyl ester, (Z,Z,Z)-	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	292.45	16.067	1.25
5	Heneicosane	C <sub>21</sub> H <sub>44</sub>	296.57	17.829	1.31
6	Cyclohexane, 1,3,5-triphenyl-	C <sub>24</sub> H <sub>24</sub>	312.44	19.157	7.01
7	2-Ethylacridine	C <sub>15</sub> H <sub>13</sub> N	207.27	23.734	1.23
8	Campesterol	C <sub>28</sub> H <sub>48</sub> O	400.68	24.781	2.11
9	1,2-Bis(trimethylsilyl)benzene	C <sub>12</sub> H <sub>22</sub> Si <sub>2</sub>	222.47	24.965	1.42
10	$\gamma$ -Sitosterol	C <sub>29</sub> H <sub>50</sub> O	414.70	25.434	8.78
11	$\beta$ -Amyrone	C <sub>30</sub> H <sub>48</sub> O	424.70	25.863	7.25
12	$\alpha$ -Amyrin	C <sub>30</sub> H <sub>50</sub> O	426.71	26.298	32.49
13	Olean-12-en-3-ol, acetate, (3 $\beta$ )-	C <sub>32</sub> H <sub>52</sub> O <sub>2</sub>	468.75	26.578	5.68
14	Lup-20(29)-en-3-ol, acetate, (3 $\beta$ )-	C <sub>32</sub> H <sub>52</sub> O <sub>2</sub>	468.75	27.030	25.72

**Comment [h19]:** Place all Figures and Tables at their appropriate positions in the text.



**Fig. 3:** Structures of compounds present in flower extract of *Vinca major*.

$\gamma$ -Sitosterol, a moderately abundant compound, has been found in many plant species such as *Lippia nodiflora* and *Acacia nilotica*, with anticancer (Sundarraj *et al.*, 2012) and antidiabetic properties (Balamurugan *et al.*, 2011). 9,12,15-octadecatrienoic acid, methyl ester, (Z,Z,Z)- and pentadecanoic acid, 14-methyl-, methyl ester are fatty acid methyl esters, which have been found in numerous plant species such as *Cannabis sativa*, *Ageratum conyzoides* and *Coronopus didymus* (Banaras *et al.*, 2021; Javaid *et al.*, 2018, 2021b). Such compounds showed antifungal and antibacterial activities (Johnson *et al.*, 2011; Bashir *et al.*, 2012). *n*-hexadecanoic acid, also known as palmitic acid, is a

highly biologically active compound, found in many plants (Javaid *et al.*, 2018; Naqvi *et al.*, 2020). It possesses anti-inflammatory, antimicrobial, mosquito larvicidal, hypocholesterolemic, antioxidant and pesticidal properties (Rahuman *et al.*, 2000; Kumar *et al.*, 2010; Aparna *et al.*, 2012; Abubakar and Majinda, 2016). Heneicosane, an alkane, isolated from *Plumbago zeylanica* showed antimicrobial activity against *Streptococcus pneumoniae* and *Aspergillus fumigatus* (Vanitha *et al.*, 2020). Likewise, 2-Ethylacridine also possesses antimicrobial and antitumor activities (Vijayakumari and Raj, 2019). Campesterol is a sterol found in plant and cholesterol lowering and anticarcinogenic properties (Choi *et al.*, 2007).

**Table 2:** Bioactivity of components of methanolic flower extract of *Vinca major*.

Sr. No.	Names of compounds	Bioactivity	Reference
1	5-Dodecene, (E)-	-	-
2	Pentadecanoic acid, 14-methyl-, methyl ester	Antifungal, Antimicrobial	Bashir <i>et al.</i> (2012)
3	<i>n</i> -Hexadecanoic acid	Antioxidant, pesticidal, anti-inflammatory, mosquito larvicide, hemolytic	Kumar <i>et al.</i> (2010); Aparna <i>et al.</i> (2012); Abubakar and Majinda (2016)
4	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	Antibacterial, antioxidant, anticancer, antipyretic, cardioprotective, antiarthritic neural function, Antiandrogenic,	Johnson <i>et al.</i> (2011); Akpuaka <i>et al.</i> (2013)
5	Heneicosane	Antimicrobial	Vanitha <i>et al.</i> (2020)
6	Cyclohexane, 1,3,5-triphenyl-	-	-
7	2-Ethylacridine	Antimicrobial, antitumor	Vijayakumari and Raj (2019)
8	Campesterol	Anti-cholesterol, anticarcinogenic	-
9	1,2-Bis(trimethylsilyl)benzene	-	-
10	$\gamma$ -Sitosterol	Anticancer, antidiabetic	Balamurugan <i>et al.</i> (2011); Sundarraj <i>et al.</i> (2012)
11	$\beta$ -Amyrone	Anti-inflammatory	de Almeida <i>et al.</i> (2015)
12	$\alpha$ -Amyrin	Antimicrobial Anti-inflammatory, antioxidant, inhibitor of human oxidosqualene cyclase	Singh <i>et al.</i> (2002); Chen <i>et al.</i> (2017); Cardoso <i>et al.</i> (2020)
13	Olean-12-en-3-ol, acetate, (3 $\beta$ )-	Antibacterial, anti-inflammatory	Hichri <i>et al.</i> (2003); Akihisa <i>et al.</i> , (2010)
14	Lup-20(29)-en-3-ol, acetate, (3 $\beta$ )-	Anticancer, anti-inflammatory, antituberculosis, antimalarial, antimicrobial, antinociceptive	Prachayasittikul <i>et al.</i> (2010); Chen <i>et al.</i> (2012); Suwito <i>et al.</i> (2016)

## Conclusion

Flower of *V. major* is a rich source of bioactive substances especially *n*-hexadecanoic acid;  $\alpha$ -amyrin, lupeol acetate,  $\gamma$ -sitosterol; heneicosane and fatty acid methyl esters with a number of biological activities including anticancer and antidiabetic.

## REFERENCES

- Abubakar, M.N. and R.R.T. Majinda. 2016. GC-MS Analysis and preliminary antimicrobial activity of *Albizia adianthifolia* (Schumach) and *Pterocarpus angolensis*. Medicines, 3: Article 3.
- Ajaib, M., S.K. Haider, A. Zikrea and M.F. Siddiqui. 2014. Ethnobotanical studies of herbs of Agra valley Parachinar, Upper Kurram Agency, Pakistan. *Int. J. Biol. Biotechnol.*, 11: 71-83.
- Akihisa, T., N. Kojima, T. Kikuchi, K. Yasukawa, H. Tokuda, E. Masters, A. Manosroi and J. Manosroi. 2010. Anti-inflammatory and chemopreventive effects of triterpene cinnamates and acetates from shea fat. *J. Oleo Sci.*, 59: 273-280.
- Akpuaka, A., M.M Ekwenchi, D.A. Dashak and A. Dildar. 2013. Biological activities of characterized isolates of *n*-hexane extract of *Azadirachta indica* A. Juss (Neem) leaves. *Nat. Sci.*, 11(5):142-145.
- Aparna, V., D. Vijayan, P. Mandal and P. Karthe. 2012. Anti-inflammatory property of *n*-hexadecanoic acid: Structural evidence and kinetic assessment. *Chem. Biol. Drug Des.*, 80: 434-439.
- Arora, R., A.K. Mathur, A. Mathur and C.M. Govil. 2010. Biotechnology of Himalayan *Vinca major* and *V. minor*. In: Medicinal Plant Biotechnology, CABI Publishers UK. pp. 207-221.
- Balamurugan, R., V. Duraipandiyan and S. Ignacimuthu. 2011. Antidiabetic activity of  $\gamma$ -sitosterol isolated from *Lippia nodiflora* L. in streptozotocin induced diabetic rats. *Eur. J. Pharmacol.*, 667: 410-418.
- Banaras, S., A. Javaid and I.H. Khan. 2021. Bioassays guided fractionation of *Ageratum conyzoides* extract for the identification of natural antifungal compounds against *Macrophomina phaseolina*. *Int. J. Agric. Biol.*, 25(4): 761-767.
- Bashir, A., K. Ibrar, B. Shumaila and S. Azam. 2012. Chemical composition and antifungal, phytotoxic, brine shrimp cytotoxicity, insecticidal, and antibacterial activities of the essential oils of *Acacia modesta*. *J. Med. Plants Res.*, 6: 4653-4659.
- Cardoso, B.K., H.L.M. de Oliveira, U.Z. Melo, C.M.M. Fernandez, C.F.A.A. Campo and J.E. Goncalves. 2020. Antioxidant activity of  $\alpha$ - and  $\beta$ -amyrin isolated from *Myrcianthes pungens* leaves. *Nat. Prod. Res.*, 34: 1777- 1781.
- Chen, Y.F., C. Ching, T.S. Wu, C.R. Wu, W.T. Hsieh and H.Y. Tsai. 2012. *Balanophora spicata* and lupeol acetate possess antinociceptive and anti-inflammatory activities *in vivo* and *in vitro*. *Evid. Based Complement. Altern. Med.*, 2012: Article 371273.
- Chen, D., F. Xu, P. Zhang, J. Deng, H. Sun, X. Wen and J. Liu. 2017. Practical synthesis of  $\alpha$ -amyrin,  $\beta$ -amyrin, and lupeol: The potential natural inhibitors of human

**Comment [h20]:** 1-References should be strictly according to PJWSR format. 2- References should be in alphabetical order. 3-References of the same first author should be in chronological order. However, references with a author should be written first.

**Comment [h21]:** Follow this format for writing names of authors and year.

**Comment [h22]:** Write name of a journal in standard abbreviated form.

**Comment [h23]:** Follow this format for writing for writing volume and page numbers.

**Comment [h24]:** For a Book Chapter, write name of chapter, name of book, publisher's name, editors and page numbers.

**Comment [h25]:** For online journals without page numbers.

oxidosqualene cyclase. Arch. Pharm. (Weinheim), 350: doi: 10.1002/ardp.201700178.

- Choi, J-M., E-O. Lee, H-J. Lee and S-H. Kim. 2007. Identification of campesterol from *Chrysanthemum coronarium* L. and its antiangiogenic activities. *Phytother. Res.*, 21: 954-959.
- Comfort, M.I., D. Majesty, A. Eze, N. Kelechi, E. Ahamefula, N. Ijeoma and O. Emmanuel. 2019. Effect of ethanolic leaf extract of *Vinca major* L. on biochemical parameters and glucose level of alloxan induced diabetic rats. *Afr. J. Biotechnol.*, 18: 1054-1068.
- Dar, R.A., M. Shahnawaz and P.H. Qazi. 2017. General overview of medicinal plants: A review. *J. Phytopharmacol.*, 6: 349-351.
- de Almeida, P.D.O., A.P.A. Boleti, A.L. Rüdiger, G.A. Lourenço, V.F. da V. Junior and E.S. Lima. 2015. Anti-inflammatory activity of triterpenes isolated from *Protium paniculatum* oil-resins. *Evid. Based Complement. Altern. Med.*, 2015: Article 293768.
- Ferdosi, M.F.H., I.H. Khan, A. Javaid, T. Sattar and A. Munir. 2020. Identification of antimicrobial constituents in essential oil of *Paulownia fortunei* flowers. *Mycopath*, 18(2): 53-57.
- Ferdosi, M.F.H., I.H. Khan, A. Javaid, M. Nadeem and A. Munir. 2021a. Biochemical profile of *Calotropis procera* flowers. *Pak. J. Weed Sci. Res.* 27(3): 341-349.
- Ferdosi, M.F.H., I.H. Khan, A. Javaid and M.F.A. Fardosi. 2021b. GC-MS examination of methanolic extract of *Cirsium arvense* flower. *Pak. J. Weed Sci. Res.*, 27(2): 173-180.
- Fernandes, C.P., A.L. Correa, J.F.R. Lob, O.P. Caramel, F.B. de Almeida and L. Rocha. 2013. Triterpene esters and biological activities from edible fruits of *Manilkara subsericea* (Mart.) Dubard, Sapotaceae. *Biomed Res. Int.*, 2013: Article 280810.
- González-Burgos, E. and M.P. Gómez-Serranillos. 2021. Vinca alkaloids as chemotherapeutic agents against breast cancer. Discovery and development of anti-breast cancer agents from natural products. In: *Natural Product Drug Discovery*, G. Brahmachari (Ed.), Elsevier. pp. 69-101.
- Hichri, F., H.B. Jannet, J. Cheriaa, S. Jegham and Z. Mighri. 2003. Antibacterial activities of a few prepared derivatives of oleanolic acid and of other natural triterpenic compounds. *C. R. Chim.*, 6(4):473-483.
- Jamshidi-Kia, F., Z. Lorigooini and H. Amini-Khoei. 2018. Medicinal plants: Past history and future perspective. *J. Herbmed. Pharmacol.*, 7: 1-7.
- Javaid, A., U. Latif, N. Akhtar, D. Ahmed and S. Perveen. 2018. Molecular characterization of *Fusarium moniliforme* and its management by methanolic extract of *Coronopus didymus*. *Pak. J. Bot.* 50(5): 2069-2075.
- Javaid, N., M.H. Shah, I.H. Khan, A. Javaid and S.M. Waleed. 2020. Herbicidal activity of *Ageratum conyzoides* against parthenium. *Pak. J. Weed Sci. Res.*, 26(2):137-146.
- Javaid, A., S.F. Naqvi and I.H. Khan. 2021a. Ethyl acetate extract of *Chenopodium murale* root, a source of bioactive compounds. *Pak. J. Weed Sci. Res.*, 27(1): 93-100.
- Javaid, A., I.H. Khan and M.F.H. Ferdosi. 2021b. Bioactive constituents of wild *Cannabis sativa* roots from Pakistan. *Pak. J. Weed Sci. Res.*, 27(3): 359-368.

Comment [h26]: For an article in press/ahead of publication/ Online First



- Javed, S., Z. Mahmood, K.M. Khan, S.D. Sarker, A. Javaid, I.H. Khan and A. Shoaib (2021). Lupeol acetate as a potent antifungal compound against opportunistic human and phytopathogenic mold *Macrophomina phaseolina*. *Sci. Rep.*, 11: 8417.
- Johnson, M., Y. Mariswamy and W.F. Gnaraj. 2011. Chromatographic finger print analysis of steroids in *Aerva lanasa* L. by HPTLC technique. *Asian Pal. J. Trop. Biomed.*, 1: 428-433.
- Khan, I.H. and A. Javaid. 2019. Antifungal, antibacterial and antioxidant components of ethyl acetate extract of quinoa stem. *Plant Prot.*, 3(3): 125-130.
- Khan, I.H. and A. Javaid. 2020. Anticancer, antimicrobial and antioxidant compounds of quinoa inflorescence. *Adv. Life Sci.*, 8(1): 68-72.
- Khan, I.H. and A. Javaid. 2021. Identification of biologically important compounds in neem leaves through GC-MS analysis. *Jordan J. Pharm. Sci.*, 14(3): 359-366.
- Kumar, P.P., S. Kumaravel and C. Lalitha. 2010. Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. *Afr. J. Biochem. Res.*, 4: 191-195.
- Manandhar, S., S. Luitel and R.K. Dahal. 2019. *In vitro* antimicrobial activity of some medicinal plants against human pathogenic bacteria. *J. Trop. Med.*, 2019: Article 1895340.
- Naqvi, S.F., I.H. Khan and A. Javaid. 2020. Hexane soluble bioactive components of *Chenopodium murale* stem. *Pak. J. Weed Sci. Res.*, 26(4): 425-432.
- Nogueira, A.O., Y.I.S. Oliveira, B.L. Adjafre, M.E.A de Moraes and G.F. Aragao. 2019. Pharmacological effects of the isomeric mixture of alpha and beta amyryn from *Protium heptaphyllum*: a literature review. *Fundam. Clin. Pharmacol.*, 33: 4-12.
- Pinto H.S.A., L.M.S. Pinto, G.M.A. Cunha, M.H. Chaves, F.A. Santos and V.S. Rao. 2019. Anti-inflammatory effect of  $\alpha$ ,  $\beta$ -amyryn, a pentacyclic triterpene from *Protium heptaphyllum* in rat model of acute periodontitis. *Inflammopharmacology*, 16: 48-52.
- Prachayasittikul, S., P. Saraban, R. Cherdtrakulkiat, S. Ruchirawat and V. Prachayasittikul. 2010. New bioactive triterpenoids and antimalarial activity of *Diospyros rubra* Lec. *EXCLI J.*, 9: 1-10.
- Rahuman, A.A., G. Gopalakrishnan, B.S. Ghouse, S. Arumugam and B. Himalayan. 2000. Effect of *Feronia limonia* on mosquito larvae. *Fitoterapia*, 71: 553-555.
- Rajput, M.S., V. Nair, A. Chauhan, H. Jawanjal and V. Dange. 2011. Evaluation of antidiarrheal activity of aerial parts of *Vinca major* in experimental animals. *Middle-East J. Sci. Res.*, 7: 784-788.
- Salmerón-Manzano, E., J.A. Garrido-Cardenas and F. Manzano-Agugliaro. 2020. Worldwide research trends on medicinal plants. *Int. J. Environ. Res. Public Health*, 17: Article ID 3376.
- Saravanan, M., P. Senthilkumar, K. Kalimuthu, V. Chinnadurai, S. Vasantharaj and A. Pugazhendhi. 2018. Phytochemical and pharmacological profiling of *Turnera subulata* Sm., a vital medicinal herb. *Ind. Crop. Prod.*, 124: 822-833.
- Shinwari, Z.K. 2010. Medicinal plants research in Pakistan. *J. Med. Plant Res.*, 4: 161-176.
- Singh, B., P.M. Sahu and M.K. Sharma. 2002. Anti-inflammatory and antimicrobial activities of triterpenoids from *Strobilanthes callosus* Nees. *Phytomedicine*, 9: 355-359.

Comment [h27]: Write full name of a journal if it is composed of a single word.

- Singh, S., J. Gupta and S.S. Kanwar. 2014. Antilipase, antiproliferatic and antiradical activities of methanolic extracts of *Vinca major*. J. Pharm. Phytochem., 3: 53-64.
- Sundarraaj, S., R. Thangam, V. Sreevani, K. Kaveri, P. Gunasekaran, S. Achiraman and S. Kannan. 2012.  $\gamma$ -Sitosterol from *Acacia nilotica* L. induces G2/M cell cycle arrest and apoptosis through c-Myc suppression in MCF-7 and A549 cells. J. Ethnopharmacol., 141: 803-809.
- Suwito, H., W.L. Heffen, H. Cahyana, and W.P. Suwarso. 2016. Isolation, transformation, anticancer, and apoptosis activity of lupeyl acetate from *Artocarpus integrus*. AIP Conf. Proc., 1718: 080004.
- Tungmunnithum, D., A. Thongboonyou, A. Pholboon and A. Yangsabai. 2018. Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview. Medicines, 5: Article ID 93.
- Ullah, N. 2017. Medicinal Plants of Pakistan: Challenges and Opportunities. Int. J. Complement. Alternat. Med., 6: Article 00193.
- Uritu, C.M., C.T. Mihai, G.D. Stanciu, G. Dodi, T. Alexa-Stratulat, A. Luca and B.I. Tamba. 2018. Medicinal plants of the family Lamiaceae in pain therapy: A review. Pain Res. Manag., 2018: Article ID 7801543.
- Vanitha, V., S. Vijayakumar, M. Nilavukkarasi, V.N. Punitha, E. Vidhya and P.K. Praseetha. 2020. Heneicosane-A novel microbicidal bioactive alkane identified from *Plumbago zeylanica* L. Ind. Crops Prod., 154: 112748.
- Vijayakumari, J. and T.L.S. Raj. 2019. GC- MS analysis of secondary metabolites from acetone and chloroform extract of *Dicranopteris linearis* (Burm. F.) Underw. Int. Res. J. Biol. Sci., 8: 39-43.
- Vishwakarma, R. and V. Prajapati. 2019. Drug of Vinca: Used as a anticancer agent". J. Pharmacol. Toxicol., 1: 40-44.
- Wei, Q. and R. Liu. 2021. Constituents of essential oil from *Vinca major* var. *variegata* and its antibacterial activity. Chem. Nat. Compd., 57: 965-967.